arXiv:0811.1548v2 [gr-qc] 12 Oct 2009

Thermodynamics of tidal charged black holes

László Árpád Gergely[1,2], Narit Pidokrajt[3], Sergei Winitzki[4]

[1] *Department of Theoretical Physics, University of Szeged,*

Tisza Lajos krt 84-86, Szeged 6720, Hungary

[2] *Department of Experimental Physics,*

University of Szeged, Dóm Tér 9, Szeged 6720, Hungary

[3] *Department of Physics, Stockholm University, 106 91 Stockholm, Sweden*

[4] *Arnold Sommerfeld Center for Theoretical Physics,*

Department of Physics, Ludwig-Maximilians University,

Theresienstr. 37, 80333 Munich, Germany

(Dated: October 12, 2009)

Abstract

We study thermodynamics and thermodynamic geometries of tidal charged black holes in four dimensions. Such black holes are spherically symmetric vacuum solutions of the effective Einstein equations on the brane and are characterized by the mass m and (generalizing their general relativistic counterparts) by a second parameter, the tidal charge q. The latter is an imprint of the Weyl curvature of the 5-dimensional space-time, in which the brane is embedded. The heat capacity of the tidal charged black hole diverges on a set of measure zero of the parameter space. However, there is no indication of phase transition at those points based on both the Poincaré stability and the thermodynamic geometry analyses, similarly to the Reissner-Nordström black hole. The uncovered thermodynamic geometries of the tidal charged black holes are as follows: the Weinhold metric is flat, the Ruppeiner metric has a positive Ricci curvature. The state space is conformal to the right half of the interior of the future light cone in a Minkowski plane.

I. INTRODUCTION

Thermodynamics of black holes (BHs) was formulated almost 40 years ago by Bardeen, Carter and Hawking [1]. The BH entropy (also known as Bekenstein-Hawking entropy) is known to be proportional to the area of the event horizon. Without using natural units, it reads

$$S = \frac{c^3 k_B A}{4G\hbar} \ , \tag{1}$$

where c, k_B, G and $\hbar$ are speed of light, Boltzmann's constant, Newton's gravitational constant and the reduced Planck's constant, respectively and A represents the area of the horizon.

Despite the fact that BH thermodynamics is well established, there is no controlled calculation of BH entropy based on standard statistical mechanics which associates entropy with a large number of microstates. The first paper on the BH's microstate counting appeared in 1996 by Strominger and Vafa [2] when they were able to calculate the Bekenstein-Hawking entropy of a five-dimensional extremal BH in the framework of string theory, and ever since there have been a growing number of papers on this topic.

An alternative way to study BH thermodynamics phenomenologically is by using a certain pseudo-Riemannian geometry defined on the thermodynamic state space (the parameter space). On such a space, one defines the "thermodynamical" metric known as the Ruppeiner metric. This metric is defined as the negative of the Hessian of the entropy S with respect to the mass m and other extensive parameters q^i, including electric charge and angular momentum:

$$g^R_{ab} = -\partial_a \partial_b S(m, q^i) \ . \tag{2}$$

The first paper on the subject is by Ruppeiner [3]; a review on the application to various thermodynamic systems is given in [4]. Recently the Ruppeiner geometry of anyon gas has also been worked out [5].

Mathematically the Ruppeiner geometry is one particular type of information geometry [6]. Ruppeiner originally developed his theory in the context of thermodynamic fluctuation theory, for systems in canonical ensembles. Many black holes have *negative* specific heats and are described microcanonically. For systems with non-interacting underlying statistical models, such as the ideal gas, the Ruppeiner geometry is flat [3]. Singularities in the curvature of the Ruppeiner metric signal thermodynamic instabilities of the system in question.

The most physically significant result as reported in [7] is that the Myers-Perry Kerr BH thermodynamic instability is detectable in the Ruppeiner geometry. More precisely, the curvature singularities of the Ruppeiner metric signal the onset of thermodynamic instabilities of the Myers-Perry Kerr BHs in dimensions $D > 5$, found earlier by Emparan and Myers [8].

In [9] it was shown that the Ruppeiner metric for a two-dimensional thermodynamic state space is flat, provided the entropy takes the power-law form $S = m^p f(Q/m)$ for any $p \neq 1$, where Q represents a conserved charge (the angular momentum in the case of the spinning Kerr BH). This small theorem is useful when one wants to quickly rule out thermodynamic curvature singularites of the system. Certain BHs do have flat Ruppeiner metric, *e.g.* Reissner-Nordström (RN), BTZ, and 4-dimensional Einstein-Maxwell-dilaton BHs. Well known examples of BHs with non-flat Ruppeiner geometry include Kerr, Kerr-Newman, and Reissner-Nordström-AdS BHs [10]. Recent papers on applications of the Ruppeiner geometry to black hole thermodynamics are listed in [11]-[19]. To date thermodynamic geometry is accepted as one of the standard tools used in investigating whether instability or critical phenomena is present in a given thermodynamic sytem.

An alternative geometric approach to thermodynamics is given by the Weinhold metric [20], which is defined as the Hessian of the mass (internal energy) with respect to entropy and other extensive parameters such as charge, angular momentum, etc.

$$g^W_{ab} = \partial_a \partial_b m(S, q^i) \ . \tag{3}$$

The Weinhold metric is equal to the conformally rescaled Ruppeiner metric,

$$g^W = T g^R \ , \tag{4}$$

where it is understood that both metrics have been transformed to the same set of coordinates. The conformal factor T is the temperature,

$$T = \partial_S m = \frac{1}{\partial_m S} \ . \tag{5}$$

The relationship (4) has been derived previously using involved thermodynamical arguments [21]. We present in the Appendix a generic and simple proof of the statement (4).

Black hole solutions arise not only in general relativity and string theory, but also in brane-world gravity models. There are many brane-world scenarios, but in the simplest gravity evolves in a curved 5D space-time (the bulk), which contains a temporal 4D hypersurface (the brane), on which all the field of the standard model are localized. Gravitational dynamics on the brane is governed by an effective Einstein equation [22]-[24].

The most well-known brane BH is the spherically symmetric vacuum *tidal* charged BH, derived in [25]:

$$ds^2 = -f\left(r\right) dt^2 + f^{-1}\left(r\right) dr^2 + r^2 \left(d\theta^2 + \sin^2\theta d\varphi^2\right) \ . \tag{6}$$

The metric function f is given as

$$f\left(r\right)=1-\frac{2m}{r}+\frac{q}{r^{2}}\,. \tag{7}$$

Such BHs are characterized by two parameters: their mass m and tidal charge q. The latter arises from the Weyl curvature of the 5D space-time into which the brane is embedded (more exactly, from its "electric" part as computed with respect to the brane normal).

Formally the metric (6) agrees with the Reissner-Nordström solution of a spherically symmetric Einstein-Maxwell system in general relativity, provided we replace the tidal charge q by the square of the *electric* charge Q. Thus $q=Q^{2}$ is always positive, when the metric (6) describes the spherically symmetric exterior of an electrically charged object in general relativity. By contrast, in brane-world theories the metric (6) allows for any sign of q. A positive tidal charge weakens the gravitational field of the black hole in precisely the same way the electric charge of the Reissner-Nordström black hole does. A negative tidal charge, however, strengthens the gravitational field, contributing to the localization of gravity on the brane.

The structure of the tidal charged BH in the case $q>0$ is in full analogy with the general relativistic Reissner-Nordström solution[1]. For $0<q<m^{2}$ it describes tidal charged BHs with two horizons, located at $r_{\pm}=m\pm\sqrt{m^{2}-q}$, both below the Schwarzschild radius. For $q=m^{2}$ the two horizons coincide at $r_{e}=m$ (this is the analogue of the extremal Reissner-Nordström black hole). Finally there is a new possibility forbidden in general relativity due to physical considerations on the smallness of the electric charge. This is $q>m^{2}$ for which the metric (6) describes a naked singularity. Such a situation can arise whenever the mass m of the brane object is small enough, compared to the effect of the Weyl curvature expressed as a tidal charge. If we assume that the tidal charge q is essentially a global property of the brane, then the latter can contain many BHs of mass $m\geq\sqrt{q}$ and several naked singularities with mass $m<\sqrt{q}$, which would be an intriguing contribution to dark matter.

For any $q<0$ there is only one horizon, at $r_{+}=m+\sqrt{m^{2}+|q|}$. For these BHs, gravity is increased on the brane by the presence of the tidal charge. This again contributes towards the localization of gravity on the brane.

Work on the tidal charged BH includes the matching with an interior stellar solution, a procedure requiring a negative q [27], the study of weak deflection of light to second order in both parameters

---

[1] In making analogy with the RN BH one can also consider the Born-Infeld BHs, which is a nonlinear generalization of the RN BH [26]

[28], [29], a confrontation with solar system tests [30], and the evolution of thin accretion disks in this geometry [31].

In this paper we analyze the tidal charged BH with the geometric methods provided by the thermodynamic metrics. We start in Section II by deriving the entropy of the tidal charged BH and analyzing its mass and tidal charge dependences. These thermodynamic considerations could be useful not only per se, but also for the analysis of the possible 5D extensions of the tidal charged BH. What we have in mind here is that some years ago Emparan, Horowitz and Myers have constructed a lower-dimensional toy model with BTZ black holes on the brane, which are BTZ "black strings" in the bulk [32]. They were able to show that the four-dimensional entropy computed from the horizon area agrees (to a leading order at large mass) with the three-dimensional entropy computed from the circumference of the horizon. The two definitions of the entropy differed in the number of dimensions and in the value of the corresponding Planck masses. Therefore the result of [32] can be taken as a hint that there are thermodynamic constraints on the 5D extension of the tidal charged BH, which are provided by our present analysis.

In Section II we also compute the temperature and analyze the heat capacity at constant q. Then in Section III we present an exhaustive study of the Ruppeiner and Weinhold geometries of the tidal charged BH. Finally we discuss our results in the concluding section.

II. THERMODYNAMIC CONSIDERATIONS

A. Entropy and mass

The exterior horizon for $q > 0$ or the only horizon for $q < 0$ are both given by

$$r_+ = m + \Theta \ , \tag{8}$$

where we have introduced the shorthand notation $\Theta = \sqrt{m^2 - q}$, real for any $q \leq m^2$. The BH's entropy can be calculated using the celebrated formula (1) with geometrized units and $k_B = 1/\pi$ chosen for convenience,

$$S = \frac{A}{4\pi} = r_+^2 = (m + \Theta)^2 \ . \tag{9}$$

It is instructive to see the variation of entropy with the BH parameters. If the mass increases, the entropy will always increase (Fig. 1). By contrast, the entropy decreases with increasing q (irrespective of the sign of q) until $S = m^2$ and the minimal horizon area $A = 4\pi m^2$ are reached at



Figure 1: The entropy always increases with the mass. The plots are for negative tidal charged BH with $q = -1$ (solid line), Schwarzschild BH with $q = 0$ (dashed line) and positive charged tidal BH with $q = 1$ (dotted line). In the latter case one can see that the entropy is not defined for masses below the extremal limit $m < \sqrt{q}$.

the extremal limit $q = m^2$ (Fig. 2). This is to be expected since for $q > m^2$ the metric (6) describes a naked singularity (thus the area A is undefined).

We can also invert the entropy function to obtain the mass for the tidal charged BH as follows

$$m = \frac{\sqrt{S}}{2}\left(1 + \frac{q}{S}\right). \tag{10}$$

Invoking the first law of thermodynamics

$$dm = TdS + \psi dq \tag{11}$$

where

$$\psi = \frac{\partial m}{\partial q} = \frac{1}{2\sqrt{S}} \tag{12}$$

is the potential associated with the tidal charge.

Although the solution (6) is static, one could interpret these findings in the following sense. In order to obey the second law of thermodynamics, the mass of such a BH can slowly increase in a classical quasi-stationary process.[2] This is exactly what is happening in an accretion process. Similarly, the 5D geometry may evolve in a quasi-stationary way only so that the tidal charge decreases in time. This is a second condition that should be obeyed by the yet unknown 5D space-time admitting the tidal charged BH brane (the first condition being that it is not 5D Schwarzschild-anti de Sitter).

---

[2] For now, we disregard the Hawking radiation, which is negligible for astrophysical black holes.



Figure 2: The qualitative behaviour of entropy vs. tidal charge. The entropy decreases with increasing q. The plot is for $m = 2$.

B. Hawking temperature and heat capacity

By the first law of thermodynamics, the Hawking temperature of the BH is given by

$$T(m,q) = \partial_S m = \frac{1}{\partial_m S} = \frac{\Theta}{2\,(m+\Theta)^2} \ . \tag{13}$$

The same value $T(m,q)$ is found by computing the temperature of the Hawking radiation if one uses the well-known formula for the surface gravity of a spherically symmetric Killing horizon (see e.g. [38]).

The temperature $T(m,q)$ increases with q for $q < 0$ up to the maximal value $T = 1/(8m)$ at $q = 0$, then decreases with increasing $q > 0$ down to $T = 0$, which is reached in the extremal limit ($q = m^2$). Thus the minimal entropy belongs to $T = 0$ and the hottest BH with a given mass is for $q = 0$, representing the Schwarzschild BH.

The heat capacity of this BH at constant q is given by

$$C_q = \frac{\partial m}{\partial T} = T\frac{\partial S}{\partial T} = T\left(\frac{\partial T}{\partial S}\right)^{-1} = T\left(\frac{\partial^2 m}{\partial S^2}\right)^{-1} = \frac{-2S(S-q)}{S-3q} \ . \tag{14}$$

We can readily observe that the heat capacity diverges along $S = 3q$ or equivalently $q = 3m^2/4$. No such behaviour occurs for any $q < 0$ (for which we always have $-\infty < C_q < 0$). In (m,q)-coordinates we can express the heat capacity as

$$C_q = 2\Theta\frac{(m+\Theta)^2}{m-2\Theta} \ . \tag{15}$$

Plotted against Θ, the heat capacity is shown on Fig 3.



Figure 3: The heat capacity C_q as function of Θ for $m = 0.5$ (solid curve), $m = 1.5$ (dashed curve) and $m = 3$ (dotted curve). All these curves show a vanishing heat capacity in the extremal case (at $\Theta = 0$) and a divergent behaviour at $\Theta = m/2$ (equivalent to $q = 3m^2/4$ or $S = 3q$). Up to the Schwarzschild limit (at $\Theta = m$, marked with a bar on the graphs) the curves apply both for tidal-charged black holes and for Reissner-Norström black holes (with electric charge $Q = \sqrt{q}$). The region of the curves for $\Theta > m$ (thus $q < 0$) applies only for tidal-charged black holes.

Both the heat capacity and its first derivative

$$\frac{dC_q}{d\Theta} = \frac{2\left(m^2 - \Theta^2\right)(m + 4\Theta)}{(m - 2\Theta)^2} \tag{16}$$

become singular at $\Theta = m/2$, that is at $q = 3m^2/4$.

In the domain of its negative values the heat capacity reaches a local maximum at the Schwarzschild configuration ($\Theta = m$), as can be seen from both Eq. (16) and Fig 3.

The thermodynamical interpretation of negative heat capacity is that such a BH cannot be in a stable equilibrium with an infinite heat reservoir held at $T = T_{BH}(m, q)$. For instance, a small thermal fluctuation may transfer some heat to the BH and make the BH colder, thus making heat transfer even more efficient. This is the typical behavior of Schwarzschild BHs, which are unstable with respect to emission of Hawking radiation in empty space and can be stable only in thermal contact with a finite-volume reservoir. Since the Universe may be considered as an infinite heat reservoir having the temperature of the cosmic background radiation, these considerations may be relevant to the cosmological stability of primordial or near-extremal BHs that have very low temperature. A near-extreme black hole with tidal charge $q > 3m^2/4$ has a positive heat capacity and thus can remain in a stable equilibrium with an infinite heat reservoir at $T = T_{BH}$.

Nevertheless, Schwarzschild BHs are known to be linearly stable with respect to perturbations which are nonvanishing on the bifurcation two sphere [33].

C. Poincaré stability analysis

In this subsection we address the issue of microcanonical stability of the tidal charged BH by using the so-called Poincaré method (see *e.g.* [34, 35, 36]) which has been used to decide whether critical phenomena or change of stability occurs in BH systems by plotting a conjugacy diagram (with a conjugacy parameter, in our case the inverse temperature, plotted against a control parameter, in our case the mass) and analyze the change in the convexity/concavity of the curve around any occuring "turning point". According to this method there is a change of stability whenever the concavity/convexity of the curve changes at a turning point.

We plot the conjugacy diagram of the tidal charged BH for $q = 0.25$ on Fig 4. In the extremal limit, at $m = 0.5$, the curve goes to infinity. The Davies point where the heat capacity diverges [37] is at $m = 0.577$. As the curve has no turning point at all, according to the Poincaré method there is no change of stability in any point, in particular neither in the Davies point.

III. THERMODYNAMIC GEOMETRY

A. The Ruppeiner metric

The geometry of the tidal charged BH depends on two parameters: m and q. From the generic definition (2) we find the corresponding Ruppeiner metric as

$$ds_R^2 = \frac{1}{\Theta^3}\left[2\left(m-2\Theta\right)\left(m+\Theta\right)^2 dm^2 - 2(m^2-\Theta^2)dmdq + \frac{m}{2}dq^2\right] \ . \tag{17}$$

The Ruppeiner curvature scalar is

$$R = \frac{1}{2\Theta(m+\Theta)} \tag{18}$$

It is readily seen that the curvature scalar diverges in the extremal limit for $q > 0$, but stays regular for any $q < 0$. It is also worth to remark that at the Davies point the metric becomes degenerate (the coefficient of dm^2 vanishes). We note here that for RN BHs the situation was identical: a singularity in the heat capacity also emerged [10] when the metric became degenerate, and was not accompanied by a phase transition.



Figure 4: A conjugacy diagram (inverse temperature vs. mass) of the tidal charged BH for $q = 0.25$. The extremal limit is at $m = 0.5$ where the curve goes to infinity. There is no "turning point", so even at $m = 0.577$, the Davies point (where the heat capacity diverges), the stability holds.

B. The Weinhold metric

By passing to coordinates $(m,\ \Theta)$ in the Ruppeiner metric and using the conformal relation (4) as well as the expression for the temperature (13), we obtain the Weinhold metric explicitly as

$$ds_W^2 = T ds_R^2 = \frac{-(m+2\Theta)\,dm^2 - 2\Theta dm d\Theta + m d\Theta^2}{(m+\Theta)^2}\ . \tag{19}$$

This can be further simplified by introducing the new coordinate r_+ replacing Θ:

$$ds_W^2 = \frac{dr_+}{r_+}\left(m\frac{dr_+}{r_+} - 2dm\right)\ , \tag{20}$$

then passing to $\left(Z = \log r_+,\ W = \log\left(r_+/m^2\right)\right)$ we find

$$ds_W^2 = m dZ dW\ , \tag{21}$$

with

$$m = \exp\left(\frac{Z-W}{2}\right)\ . \tag{22}$$

In the coordinates $(U_+ = 2\exp(Z/2)\,,\ U_- = 2\exp(-W/2))$ the Weinhold metric becomes manifestly flat, $ds_W^2 = -dU_+ dU_-$. One can also introduce Minkowskian coordinates as $U_\pm = X \pm Y$, finding $ds_W^2 = -dX^2 + dY^2$. The sequence of coordinate transformations leading to this result can be summarized as

$$\begin{aligned} X &= \sqrt{r_+} + \frac{m}{\sqrt{r_+}} \ , \\ Y &= \sqrt{r_+} - \frac{m}{\sqrt{r_+}} \ . \end{aligned} \tag{23}$$

The inverse transformation is

$$\begin{aligned} 4r_+ &= (X+Y)^2 \ , \\ 4m &= X^2 - Y^2 \ . \end{aligned} \tag{24}$$

C. Direct derivation of the Weinhold metric

As a further consistency check of our calculations we express from Eq. (9) the mass m of the tidal charged BH in (S, q) coordinates, obtaining

$$m = \frac{\sqrt{S}}{2}\left(1 + \frac{q}{S}\right) \tag{25}$$

This allows to calculate the Weinhold metric directly from its definition:

$$ds_W^2 = \frac{3q - S}{8S^{5/2}} dS^2 - \frac{1}{2S^{3/2}} dS dq \ . \tag{26}$$

It can easily be checked that the above metric is flat. It is worth noting that when $S = 3q$ (where the heat capacity diverges) the Weinhold metric is degenerate. By performing a transformation to (m, Θ) coordinates, we recover the form (19) of the Weinhold metric.

D. The geodesics of the thermodynamic metrics

The thermodynamic metrics have interesting physical interpretation. A Weinhold geodesic (the path of least Weinhold length) in the parameter space corresponds to a quasistatic process that, in a well-defined sense, optimizes the “thermodynamical time,” which roughly means the time necessary for quasistatic relaxation during the process [39]. Since in the present case the Weinhold metric is flat, its geodesics are straight lines both in the Minkowskian coordinates (X, Y) and in the null coordinates $U_\pm = X \pm Y$:

$$U_+ = \alpha U_- + \beta \ , \tag{27}$$

with α, β constants.



Figure 5: A state space plot of the tidal charged BH embedded in the flat Minkowskian parameter space. Note that the thermodynamic light cone (TLC) describes $m = 0$ and the wedge fills the right half of the TLC with the TLC itself excluded. The vertical axis represents the extremal limit in which $T = 0$.

On the other hand, the "length" of a process computed in the Ruppeiner metric describes the minimum dissipation (increase of entropy) that is unavoidable in a quasistatic process [40]. The geodesic equations for the Ruppeiner metric are the simplest when written in the conformally Minkowskian null coordinates $U_{\pm}$:

$$\begin{aligned} \frac{d^2U_+}{d\lambda^2} + \frac{2U_+ - 3U_-}{U_+ \left(U_+ - U_-\right)} \left(\frac{dU_+}{d\lambda}\right)^2 &= 0 \ , \\ \frac{d^2U_-}{d\lambda^2} + \frac{1}{U_+ - U_-} \left(\frac{dU_-}{d\lambda}\right)^2 &= 0 \ , \end{aligned} \tag{28}$$

with λ an affine parameter.

E. The global structure of the Ruppeiner geometry

The expression of the temperature in the $(X,\ Y)$ coordinates is

$$T = \frac{r_+ - m}{2r_+^2} = \frac{4Y}{\left(X + Y\right)^3} \ , \tag{29}$$

which leads to the manifestly conformally flat form of the Ruppeiner metric:

$$ds_R^2 = \frac{\left(X + Y\right)^3}{4Y} \left(-dX^2 + dY^2\right) \ . \tag{30}$$

Note that the domain of the original Ruppeiner coordinates is

$$m \in (0,\ \infty) \ , \ q \in \left(-\infty,\ m^2\right) \ . \tag{31}$$

The corresponding ranges of the variables Θ, r_+ are $\Theta \geq 0,\ r_+ \geq 0$; the Minkowskian coordinates defined by (23) have the range $X > Y \geq 0$. Thus the state space is equivalent to the right half of the interior of the future light cone of a Minkowski plane, with the vertical boundary included but the light-like boundary excluded. (The light cone describes $m = 0$ states as can be seen from $4m = X^2 - Y^2$, which for $q \geq 0$ does not correspond to BH metrics.) The extremal states are located at $(X = 2\sqrt{m} > 0,\ Y = 0)$, i.e. on the positive half of the time-like coordinate axis (the vertical boundary). This can be also seen by writing the curvature scalar (18) of the Ruppeiner metric in the (X, Y) coordinates:

$$R = \frac{1}{2\left(r_+ - m\right) r_+} = \frac{4}{Y\left(X + Y\right)^3} \ . \tag{32}$$

We also remark that passing to the $(X,\ Y)$ coordinates by the transformation (24) induces a degeneracy. For each pair of coordinates $(m,\ q)$, as well as for (m, Θ) or $(m,\ r_+)$, we can associate any of the combinations $(\pm X,\ \pm Y)$, with $X,\ Y$ defined by Eq. (23). Therefore the light cone of the Minkowski plane provides a four-fold coverage of the original state space. This is similar to the introduction of the well-known Kruskal coordinates for the Schwarzschild geometry: Kruskal coordinates cover four patches in the Kruskal-Szekeres diagram, while the original coordinates cover only one patch.

IV. CONCLUDING REMARKS

We have studied the thermodynamics and thermodynamic geometries (both Ruppeiner and Weinhold) of the tidal charged brane BH (characterized by mass m and tidal charge q) in detail.

Depending on the Weyl curvature of the 5D space-time, the induced tidal charge can be either positive or negative. Whilst from the point of view of gravitational curvature the first case is a mere rewriting of the RN geometry with electric charge $Q = \sqrt{q}$, a negative tidal charge has no correspondence in general relativity. Along the well-known property that such a negative tidal charge strengthens the gravitational field of the BH and as such contributes to the localization of gravity on the brane, our analysis has explicitly shown here that $q < 0$ is also thermodynamically preferred, as it comes with a higher entropy.

Despite the fact that the space-time about the tidal charged BH with $q > 0$ is similar to that of the RN BH, their thermodynamic geometries are quite different. Most notably, the Ruppeiner geometry of the tidal charged BH is found non-flat, as opposed to the Ruppeiner metric for the RN BH, which is flat [10].

While the thermodynamic state space of the RN BH is a Rindler wedge embedded in a Minkowski parameter space, the state space of the tidal charged BH is the right half of the interior of the future light cone of the Minkowski plane, with the vertical boundary included but the light-like boundary excluded. The light cone of the Minkowski plane provides a four-fold coverage of this state space, similarly to the four-fold covering of the curvature coordinates for a Schwarzschild BH by Kruskal coordinates.

As for some other BHs, the induced Ruppeiner geometry of the tidal charged BH has a physical singularity in the extremal limit and a degeneracy at the Davies point, where the heat capacity diverges (at $q = 3m^2/4$). The Poincaré method analysis shows however no critical phenomena ocurring there, similarly as for the Kerr BH [34], where despite the change of sign of the heat capacity through infinite values at the Davies point, the stability remains unchanged.

V. ACKNOWLEDGMENTS

LÁG wishes to acknowledge discussions with Shinji Mukohyama on the relation of the entropies of higher dimensional black objects and their lower dimensional sections. He was succesively supported by the Hungarian Scientific Research Fund (OTKA) grant no. 69036, a London South Bank University Research Opportunities Fund Award, the Polányi and Sun Programs of the Hungarian National Office for Research and Technology (NKTH) and Collegium Budapest. NP was supported by Doktorandtjänst of Stockholm University during the early stage of this work and is grateful for scholarships from Helge Ax:son Johnsons stiftelse and K & A Wallenbergs stiftelse, also for support from V. Källkvist during all the work on this paper. He would like to thank Roberto Emparan for insightful discussions and useful comments, and also wishes to acknowledge the hospitality of Departament de Física Fonamental at Universitat de Barcelona during his research visit.

Appendix A: THE GENERAL RELATION BETWEEN THE RUPPEINER AND WEINHOLD METRICS

In this Appendix we present a concise derivation of the generic statement that the Ruppeiner and Weinhold metrics (2) and (3) are conformally related, with the temperature (5) as the conformal factor.

Let us consider a scalar function[3] $m\left(x\right)$ of the variables $x \equiv \left\{x^a \mid a = \overline{1, n+1},\ n \in N\right\}$. We will also consider another coordinate system $y \equiv \left\{ y^c \mid c = \overline{1, n+1},\ n \in N \right\}$ in the same configuration space. A series expansion of $\delta m \equiv m(x + \delta x) - m(x)$ about an arbitrary point gives

---

[3] The function m is generic at this stage, but will be identified with the mass at the end of the Appendix.

$$\delta m\left(x\right) = \frac{\partial m}{\partial x^a}\delta x^a + \frac{1}{2}\frac{\partial^2 m}{\partial x^a \partial x^b}\delta x^a \delta x^b + \mathcal{O}\left(\delta x\right)^3 \ , \tag{A1}$$

$$\delta m\left(y\right) = \frac{\partial m}{\partial y^c}\delta y^c + \frac{1}{2}\frac{\partial^2 m}{\partial y^c \partial y^d}\delta y^c \delta y^d + \mathcal{O}\left(\delta y\right)^3 \ . \tag{A2}$$

It is understood that all derivatives are evaluated at the same point x_0, and that y_0 and δy_0 represent x_0 and δx_0 expressed in the coordinates $\{y^c\}$. In particular, we have a similar expansion

$$\delta x^a\left(y\right) = \frac{\partial x^a}{\partial y^c}\delta y^c + \frac{1}{2}\frac{\partial^2 x^a}{\partial y^c \partial y^d}\delta y^c \delta y^d + \mathcal{O}\left(\delta y\right)^3 \ . \tag{A3}$$

Let us now derive a relationship between $\partial^2 m / \partial x^a \partial x^b$ and $\partial^2 m / \partial y^c \partial y^d$. By inserting Eq. (A3) into Eq. (A1), we obtain a second expression for δm as a function of the coordinates y; that expression should coincide with Eq. (A2) order by order. By identifying the first-order and second-order contributions with the respective quantities of Eq. (A2) we obtain:

$$\frac{\partial m\left(y\right)}{\partial y^c} = \frac{\partial m\left(x\right)}{\partial x^a}\frac{\partial x^a\left(y\right)}{\partial y^c} \ , \tag{A4}$$

$$\frac{\partial^2 m\left(y\right)}{\partial y^c \partial y^d} = \frac{\partial^2 m\left(x\right)}{\partial x^a \partial x^b}\frac{\partial x^a\left(y\right)}{\partial y^c}\frac{\partial x^b\left(y\right)}{\partial y^d} + \frac{\partial m\left(x\right)}{\partial x^a}\frac{\partial^2 x^a\left(y\right)}{\partial y^c \partial y^d} \ . \tag{A5}$$

While Eq. (A4) is simply the chain rule, Eq. (A5) is the identity that will eventually yield the desired result.

In order to achieve this, we specify the coordinates as follows,

$$x = \left\{ q^1, ..., q^n, S \right\} \ , \tag{A6}$$

$$y = \left\{ q^1, ..., q^n, m \right\} \ . \tag{A7}$$

In other words, the coordinates y^i are the same as x^i except for the last coordinate y^{n+1}, which is chosen as the function $m(x)$ itself. It then follows that

$$\frac{\partial m(y)}{\partial y^i} = \delta_i^{n+1} \, , \tag{A8}$$

thus the left hand side of Eq. (A5) identically vanishes. It is also straightforward to see that

$$\frac{\partial^2 x^a\left(y\right)}{\partial y^c \partial y^d} = \frac{\partial}{\partial y^d}\left[\delta_i^a \delta_c^i + \delta_{n+1}^a \frac{\partial S\left(y\right)}{\partial y^c}\right] = \delta_{n+1}^a \frac{\partial^2 S\left(y\right)}{\partial y^c \partial y^d} \, . \tag{A9}$$

Thus Eq. (A5) is reduced to the identity

$$\frac{\partial^2 m\left(x\right)}{\partial x^a \partial x^b}\frac{\partial x^a\left(y\right)}{\partial y^c}\frac{\partial x^b\left(y\right)}{\partial y^d} = -\frac{\partial m\left(x\right)}{\partial S}\frac{\partial^2 S\left(y\right)}{\partial y^c \partial y^d} \ . \tag{A10}$$

By recalling the definitions (2), (3) and (5) of the Ruppeiner metric, Weinhold metric and temperature, respectively, by having in mind that the set of coordinates x and y are given by Eqs. (A6)-(A7), we have recovered the desired relation

$$g^{W}_{ab}\left(x\right)\frac{\partial x^{a}}{\partial y^{c}}\frac{\partial x^{b}}{\partial y^{d}}=Tg^{R}_{cd}\left(y\right)\ , \tag{A11}$$

where the left hand side represents the Weinhold metric transformed to the y coordinates.

---

[1] J. M. Bardeen, B. Carter and S.W. Hawking, Commun. Math. Phys. **31** 161 (1973).

[2] A. Strominger and C. Vafa, Phys. Lett. B **379**, 99 (1996) [arXiv:hep-th/9601029].

[3] G. Ruppeiner, Phys. Rev. A **20**, 1608 (1979).

[4] G. Ruppeiner, Rev. Mod. Phys. **67**, 605 (1995).

[5] B. Mirza and H. Mohammadzadeh, Phys. Rev. E **80**, 011132 (2009) [arXiv:0808.0241 [cond-mat]].

[6] D. A. Johnston, W. Janke and R. Kenna, Acta Phys. Polon. B **34**, 4923 (2003) [arXiv:cond-mat/0308316].

[7] J. E. Åman and N. Pidokrajt, Phys. Rev. D **73**, 024017 (2006) [arXiv:hep-th/0510139].

[8] R. Emparan and R. C. Myers, JHEP **0309**, 025 (2003) [arXiv:hep-th/0308056].

[9] J. E. Åman, I. Bengtsson and N. Pidokrajt, Gen. Rel. Grav. **38**, 1305 (2006) [arXiv:gr-qc/0601119].

[10] J. E. Åman, I. Bengtsson and N. Pidokrajt, Gen. Rel. Grav. **35**, 1733 (2003) [arXiv:gr-qc/0304015].

[11] J. E. Åman, *et al.* J. Phys. Conf. Ser. **66**, 012007 (2007) [arXiv:gr-qc/0611119].

[12] T. Sarkar, G. Sengupta and B. Nath Tiwari, JHEP **0611**, 015 (2006) [arXiv:hep-th/0606084].

[13] B. Mirza and M. Zamani-Nasab, JHEP **0706**, 059 (2007) [arXiv:0706.3450 [hep-th]].

[14] H. Quevedo, Gen. Rel. Grav. **40**, 971 (2008) [arXiv:0704.3102 [gr-qc]].

[15] G. Ruppeiner, Phys. Rev. D **78**, 024016 (2008) [arXiv:0802.1326 [gr-qc]].

[16] J. L. Alvarez, H. Quevedo and A. Sanchez, Phys. Rev. D **77**, 084004 (2008) [arXiv:0801.2279 [gr-qc]].

[17] Y. S. Myung, Y. W. Kim and Y. J. Park, Phys. Lett. B **663**, 342 (2008) [arXiv:0802.2152 [hep-th]].

[18] A. J. M. Medved, Mod. Phys. Lett. A **23**, 2149 (2008) [arXiv:0801.3497 [gr-qc]].

[19] J. Shen, R.-G. Cai, B. Wang and R.-K. Su, Int. J. Mod. Phys. A **22**, 11 (2007) [arXiv:gr-qc/0512035].

[20] F. Weinhold, J. Chem. Phys. **63**, 2479 (1975).

[21] P. Salamon, J. Nulton and E. Ihrig, J. Chem. Phys. **80**, 436 (1984).

[22] T. Shiromizu, K.I. Maeda and M. Sasaki, Phys. Rev. D **62,** 024012 (2000) [arXiv:gr-qc/9910076].

[23] L. Á. Gergely, Phys. Rev. D **68**, 124011 (2003) [arXiv:gr-qc/0308072].

[24] L. Á. Gergely, Phys. Rev. D **78**, 084006 (2008) [arXiv:0806.3857 [gr-qc]].

[25] N. Dadhich, R. Maartens, P. Papadopoulos and V. Rezania, Phys. Lett. B **487**, 1 (2000) [arXiv:hep-th/0003061].

[26] Y. S. Myung, Y. W. Kim and Y. J. Park, Phys. Rev. D **78**, 084002 (2008) [arXiv:0805.0187 [gr-qc]].

[27] C. Germani and R. Maartens, Phys. Rev. D **64**, 124010 (2001) [arXiv:hep-th/0107011].

[28] L. Á. Gergely and B. Darázs, Publ. Astron. Dept. Eötvös Univ. PADEU **17**, 213 (2006) [arXiv:astro-ph/0602427].

[29] L. Á. Gergely, Z. Keresztes and M. Dwornik, Class. Quantum Grav. **26**, 145002 (2009) [arXiv:0903.1558 [gr-qc]].

[30] C. G. Boehmer, T. Harko and F. S. N. Lobo, Class. Quant. Grav. **25**, 045015 (2008) [arXiv:0801.1375].

[31] C. S. J. Pun, Z. Kovacs and T. Harko , Phys. Rev. D **78**, 084015 (2008) [arXiv:0809.1284].

[32] R. Emparan, G. T. Horowitz and R. C. Myers, JHEP 0001 021 (2000) [arXiv:hep-th/9912135].

[33] B. S. Kay and R. M. Wald, Class. Quantum Grav. **4**, 893 (1987).

[34] G. Arcioni and E. Lozano-Tellechea, Phys. Rev. D **72**, 104021 (2005) [arXiv:hep-th/0412118].

[35] O. Kaburaki, I. Okamoto and J. Katz, Phys. Rev. **D47** (1993) 2234.

[36] J. Katz, I. Okamoto and O. Kaburaki, Class. Quant. Grav. **10**, 1323 (1993).

[37] P. C. W. Davies, Proc. Roy. Soc. Lond. A **353**, 499-521 (1977).

[38] T. Padmanabhan, Phys. Rept. **406**, 49 (2005) [arXiv:gr-qc/0311036].

[39] L. Diósi, K. Kulacsy, B. Lukács and A. Rácz, J. Chem. Phys. **105**, 11220 (1996).

[40] J. Nulton, P. Salamon, B. Andresen and Qi Anmin, J. Chem. Phys. **83**, 334 (1985).